



Figure 1
 Algorithm for PQRI, Measure 3 (See text).
 Diabetic patients must have one of the listed ICD-9 codes, meet age criteria, and have a recent encounter during the reporting period that meets the listed CPT or HCPCS codes.

If no documented blood pressure, use CPT II code, 2000F-8P.

Both a systolic and a diastolic code are required.

Performance is met if CPT II Codes (3074F or 3075F) and 3078F. Performance is not met if 3079F or 3080F or 3077F

SBP - Systolic blood pressure in mm Hg
 DBP - Diastolic blood pressure in mm Hg

***ICD-9 CODES FOR THIS MEASURE**

250.00, 250.01, 250.02, 250.03, 250.10, 250.11, 250.12, 250.13, 250.20, 250.21, 250.22, 250.23, 250.30, 250.31, 250.32, 250.33, 250.40, 250.41, 250.42, 250.43, 250.50, 250.51, 250.52, 250.53, 250.60, 250.61, 250.62, 250.63, 250.70, 250.71, 250.72, 250.73, 250.80, 250.81, 250.82, 250.83, 250.90, 250.91, 250.92, 250.93, 648.00, 648.01, 648.02, 648.03, 648.04

****CPT/HCPCS CODES FOR THIS MEASURE**

CPT

97802, 97803, 97804, 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99304, 99305, 99306, 99307, 99308, 99309, 99310, 99324, 99325, 99326, 99327, 99328, 99334, 99335, 99336, 99337, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350

HCPCS G0270, G0271